



PERSPECTIVES

The role of penicillin-binding protein 2 (PBP2) in the cephalosporin susceptibility of *Neisseria gonorrhoeae* and the need for consensus in naming of PBP2

Chun-Chen Chen, Kun-Yen Lin, Shu-Ying Li*

Research and Diagnostic Center, Centers for Disease Control, Taipei 105, Taiwan

Received 5 March 2012; received in revised form 9 March 2012; accepted 15 March 2012

Recently, there has been growing concern about the emergence and transmission of *Neisseria gonorrhoeae* multidrug-resistant clones, especially those displaying reduced susceptibility to cephalosporins, e.g., cefixime and ceftriaxone.^{1–3} In addition, increasing reports on treatment failures of oral cefixime, and later of injectable ceftriaxone, currently the recommended first-line antibiotic for gonorrhea treatment in many countries, have raised serious concerns worldwide.^{1,4} Genetic mechanisms mediating reduced cephalosporin susceptibility of *N. gonorrhoeae* is complicated, including mutations in several chromosomal genes such as *penA*, the gene that encodes the penicillin-binding protein 2 (PBP2); *porB1*, an outer membrane protein channel related to antibiotic entry; *ponA*, the penicillin-binding protein 1 (PBP1); *pilQ*, the outer membrane secretin PilQ; and *mtrR*, a repressor of the MtrC-MtrD-MtrE efflux pump.

Among them, the polymorphisms in the *penA* allele are the major contributors of resistance to cephalosporins in *N. gonorrhoeae*.^{2,4–6} PBP2 is a membrane-bound enzyme involved in cell wall biosynthesis and it is one of the targets of the β -lactam antibiotics. The mutation in

different positions of PBP2 may result in various levels of minimal inhibition concentration (MIC) increase. PBP2s with mosaic structure confer significantly higher MIC to cephalosporins than those having non-mosaic structures.^{1,2,4–6}

Several reports on the emergence of cephalosporin-resistant gonorrhea have indicated that the resistant strains are mostly clonal and contain specific PBP2 types, such as mosaic types X, XXXIV, and others.⁷ Thus, in view of the important role of PBP2 types in cephalosporin resistance, it is essential that the PBP2 types be characterized precisely and named correctly.

We reviewed and summarized the names of PBP2s used in different gonococcal studies during 2005–2011, and found several discrepancies in the naming of PBP2. Since 2005, at least 37 sequence types of PBP2 have been reported.^{1,2,4–6,8–12} For PBP2 sequence types I to XXIII (except types X and XII), no conflict in assignment of PBP2 names was found among previous studies. However, for the rest of the PBP2s, disagreements in nomenclature were found: (1) seven PBP2 sequence types have discordant names in different reports; and (2) under the PBP2 type XXIV, there are two alternatives, one is mosaic and the other is nonmosaic (see Table 1). In view of the increasing international transmission of gonococcal strains with reduced susceptibility to cephalosporins, unification in PBP2 nomenclature is needed.

* Corresponding author. Research and Diagnostic Center, Centers for Disease Control, Number 161, Kun-Yang Street, Taipei, Taiwan.
E-mail address: syl@cdc.gov.tw (S.-Y. Li).

Table 1 The names of different PBP2 sequence types used in previous studies.

PBP2 name based on ref. 4	Reference	PBP2 name based on refs. 1, 2, 5, and 6	Reference	Other name ^a	Reference
X ^{a,b}	4	XXXV*	5		
XII		XXXVI	5		
XXIV		XXIV	6		
XXV*		XXV*	2	mosaic-3	10
XXVI*		XXVI*	2	mosaic-4	10
XXVII*		XXIV*	2	mosaic-2	10
XXVIII*		XXVII*	2	PBP2 of strain 30/02	8
XXIX*		XXVIII*	1, 2	PBP2 of strain 35/02 ^c	8
XXXIII		XXIX	2	PBP2 of strain 188/03	8
XXXV*		XXXVIII*	5		
XXXVI					

^a The name of a PBP2 sequence type in its source study.

^b *indicates mosaic PBP2 types.

^c The PBP2 of strain 35/02 in refs. 1, 2, and 8 was assigned as type XXVIII, but the amino acid sequence of *penA* allele in *N. gonorrhoeae* strain 35/02 (GenBank accession number ZP_06129591) in NCBI was identical to PBP2 type X. We are not sure whether these two alleles are different.

Acknowledgments

This work was supported by grant DOH101-DC-2007 from the Centers for Disease Control, Department of Health.

References

- Unemo M, Golparian D, Nicholas R, Ohnishi M, Galloway A, Sednaoui P. High-level cefixime- and ceftriaxone-resistant *Neisseria gonorrhoeae* in France: novel *penA* mosaic allele in a successful international clone causes treatment failure. *Antimicrob Agents Chemother* 2011;**56**:1273–80.
- Ohnishi M, Watanabe Y, Ono E, Takahashi C, Oya H, Kuroki T, et al. Spread of a chromosomal cefixime-resistant *penA* gene among different *Neisseria gonorrhoeae* lineages. *Antimicrob Agents Chemother* 2010;**54**:1060–7.
- Li S-Y, Wong W-W. Antimicrobial resistance and molecular epidemiology of *Neisseria gonorrhoeae* in Taiwan. *J Formos Med Assoc* 2009;**108**:681–2.
- Ohnishi M, Golparian D, Shimuta K, Saika T, Hoshina S, Iwasaku K, et al. Is *Neisseria gonorrhoeae* initiating a future era of untreatable gonorrhea?: detailed characterization of the first strain with high-level resistance to ceftriaxone. *Antimicrob Agents Chemother* 2011;**55**:3538–45.
- Allen VG, Farrell DJ, Rebbapragada A, Tan J, Tijet N, Perusini SJ, et al. Molecular analysis of antimicrobial resistance mechanisms in *Neisseria gonorrhoeae* isolates from Ontario, Canada. *Antimicrob Agents Chemother* 2011;**55**:703–12.
- Lee SG, Lee H, Jeong SH, Yong D, Chung GT, Lee YS, et al. Various *penA* mutations together with *mtrR*, *porB* and *ponA* mutations in *Neisseria gonorrhoeae* isolates with reduced susceptibility to cefixime or ceftriaxone. *J Antimicrob Chemother* 2010;**65**:669–75.
- Li S-Y. Global transmission of multiple-drug resistant *Neisseria gonorrhoeae* strains refractive to cephalosporin treatment. *J Formos Med Assoc* 2012;**111**:463–4.
- Lindberg R, Fredlund H, Nicholas R, Unemo M. *Neisseria gonorrhoeae* isolates with reduced susceptibility to cefixime and ceftriaxone: association with genetic polymorphisms in *penA*, *mtrR*, *porB1b*, and *ponA*. *Antimicrob Agents Chemother* 2007;**51**:2117–22.
- Pandori M, Barry PM, Wu A, Ren A, Whittington WL, Liska S, et al. Mosaic penicillin-binding protein 2 in *Neisseria gonorrhoeae* isolates collected in 2008 in San Francisco, California. *Antimicrob Agents Chemother* 2009;**53**:4032–4.
- Takahata S, Senju N, Osaki Y, Yoshida T, Ida T. Amino acid substitutions in mosaic penicillin-binding protein 2 associated with reduced susceptibility to cefixime in clinical isolates of *Neisseria gonorrhoeae*. *Antimicrob Agents Chemother* 2006;**50**:3638–45.
- Ito M, Deguchi T, Mizutani KS, Yasuda M, Yokoi S, Ito S, et al. Emergence and spread of *Neisseria gonorrhoeae* clinical isolates harboring mosaic-like structure of penicillin-binding protein 2 in central Japan. *Antimicrob Agents Chemother* 2005;**49**:137–43.
- Whiley DM, Limnios EA, Ray S, Sloots TP, Tapsall JW. Diversity of *penA* alterations and subtypes in *Neisseria gonorrhoeae* strains from Sydney, Australia, that are less susceptible to ceftriaxone. *Antimicrob Agents Chemother* 2007;**51**:3111–6.